

## Technical Data Sheet

Product Name	BAST Fe
Description	A liquid concentrated fertilizer with a high ferrous content.
Appearance	dark brown liquid
Solubility	100%
Typical pH	4.5-6.5
Typical Density	1.27±0.05 g/cm <sup>3</sup>
Typical Analysis	70 g/l - ferrous (Fe)
Packing	1 l, 5 l, 10 l

## APPLICATION RATES AND TIMINGS

CROPS	BAST Fe Foliar application
<b>FRUIT TREES</b>	1.5-2.0 l/ha at flowering stage up to fruiting (14 - 28 day intervals). Water rate: 500-1000 l/ha.
<b>VEGETABLES (FIELD GROWN)</b>	1.0-1.5 l/ha from 6-10 true leaf stage up to fruiting at 14 to 28 day intervals. Water rate: 200-300 l/ha.
<b>GRAPEVINES</b>	1.5-2.0 l/ha at flowering stage up to fruiting at 14 to 28 day intervals. Water rate: 500-1000 l/ha.
<b>BRASSICAS</b>	1-1.5 l/ha. From 2-3 true leaf stage up to the end of the head formation stage. Application with an interval of 14-28 days. Water rate: 200-4000 l/ha.
<b>OTHER CROPS</b>	0.5-1.5 l/ha at visible ferrous (Fe) deficiency symptoms or according to the results of the analysis. For moderate to severe deficiency, a repeat application may be necessary 10 to 14 days later.